



Resistance of plant-plant networks to biodiversity loss and secondary extinctions following simulated environmental changes

Losapio, Gianalberto ; Schöb, Christian

Abstract: 1. Plant interactions are fundamental processes for structuring plant communities and are an important mechanism governing the response of plant species and communities to environmental changes. Thus, understanding the role played by the interaction network in modulating the impact of environmental changes on plant community composition and diversity is crucial. Here, we aimed to develop a new analytical and conceptual framework to evaluate the responses of plant communities to environmental changes. 2. This framework uses functional traits as sensitivity measures for simulated environmental changes and assesses the consequences of microhabitat loss. We show here its application to an alpine plant community where we recorded functional traits (specific leaf area [SLA] and leaf dry matter content [LDMC]) of all plants associated with three foundation species or the surrounding open areas. We then simulated primary species loss based on different scenarios of environmental change and explored community persistence to the loss of foundation species. 3. Generally, plant community responses differed among environmental change scenarios. In a scenario of increasing drought alone (i.e. species with lower LDMC were lost first) or increasing drought with increasing temperature (i.e. species with lower LDMC and higher SLA were lost first), the plant community resisted because drought-tolerant foundation species tolerated those deteriorating conditions. However, in scenarios with increasing nitrogen input (i.e. species having lower SLA were lost earlier), foundation species accelerated species loss due to their early primary extinctions and the corresponding secondary extinctions of species associated to their microhabitat. 4. The resistance of a plant community depends on the driver of environmental change, meaning that the prediction of the fate of this system is depending on the knowledge of the main driver of environmental change. Our framework provides a mechanistic understanding of an ecosystem response to such environmental changes thanks to the integration of biology-informed criteria of species sensitivities to environmental factors into a network of interacting species.

DOI: <https://doi.org/10.1111/1365-2435.12839>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-134670>

Journal Article

Accepted Version

Originally published at:

Losapio, Gianalberto; Schöb, Christian (2017). Resistance of plant-plant networks to biodiversity loss and secondary extinctions following simulated environmental changes. *Functional Ecology*, 31(5):1145-1152.

DOI: <https://doi.org/10.1111/1365-2435.12839>

DR GIANALBERTO LOSAPIO (Orcid ID : 0000-0001-7589-8706)
MR. CHRISTIAN SCHÖB (Orcid ID : 0000-0003-4472-2286)

Received Date : 27-Jul-2016
Revised Date : 20-Jan-2017
Accepted Date : 26-Jan-2017
Article type : Standard Paper
Editor: Gaku Kudo
Section: Community Ecology

Resistance of plant–plant networks to biodiversity loss and secondary extinctions following simulated environmental changes

Gianalberto Losapio* & Christian Schöb

Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

Correspondence author: E-mail: gianalberto.losapio@ieu.uzh.ch

Running headline: Networks' responses to environmental changes

Summary

1. Plant interactions are fundamental processes for structuring plant communities and are an important mechanism governing the response of plant species and communities to environmental changes. Thus, understanding the role played by the interaction network in modulating the impact of environmental changes on plant community composition and diversity is crucial. Here, we aimed to develop a new analytical and conceptual framework to

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1365-2435.12839

This article is protected by copyright. All rights reserved.

evaluate the responses of plant communities to environmental changes.

2. This framework uses functional traits as sensitivity measures for simulated environmental changes and assesses the consequences of microhabitat loss. We show here its application to an alpine plant community where we recorded functional traits (specific leaf area [SLA] and leaf dry matter content [LDMC]) of all plants associated with three foundation species or the surrounding open areas. We then simulated primary species loss based on different scenarios of environmental change and explored community persistence to the loss of foundation species.

3. Generally, plant community responses differed among environmental change scenarios. In a scenario of increasing drought alone (i.e. species with lower LDMC were lost first) or increasing drought with increasing temperature (i.e. species with lower LDMC and higher SLA were lost first), the plant community resisted because drought-tolerant foundation species tolerated those deteriorating conditions. However, in scenarios with increasing nitrogen input (i.e. species having lower SLA were lost earlier), foundation species accelerated species loss due to their early primary extinctions and the corresponding secondary extinctions of species associated to their microhabitat.

4. The resistance of a plant community depends on the driver of environmental change, meaning that the prediction of the fate of this system is depending on the knowledge of the main driver of environmental change. Our framework provides a mechanistic understanding of an ecosystem response to such environmental changes thanks to the integration of biology-informed criteria of species sensitivities to environmental factors into a network of interacting species.

Keywords: biodiversity loss, ecological networks, drought, facilitation, foundation species, nitrogen deposition, plant–plant interactions, trait-based extinction model.

This article is protected by copyright. All rights reserved.

Introduction

There is evidence that global environmental changes are affecting the stability of ecosystems (Hautier *et al.* 2015) causing a worldwide decline in biodiversity (Cardinale *et al.* 2012) with unprecedented high extinction rates (Pimm *et al.* 2014). However, the consequences of environmental changes for biotic interactions are much less known (Bascompte & Jordano 2014; Michalet *et al.* 2014), despite the importance of species interactions for mediating species tolerance and community persistence under environmental changes (Ives & Cardinale 2004; Fortuna & Bascompte 2006; Memmott *et al.* 2007; Saavedra *et al.* 2013).

Plant–plant interactions are fundamental processes for structuring plant communities (Klanderud 2005; Michalet *et al.* 2015). In alpine ecosystems, where harsh environmental conditions prevail (e.g. low temperature, poor soil), stress-tolerant foundation species (i.e. species that influence community structure and ecosystem processes, *sensu* Ellison *et al.* 2005) often promote species survival and growth, and community-level diversity, through creation of unique microhabitats (Cavieres *et al.* 2014). These effects of foundation species have been attributed to facilitative effects of foundation species on their co-occurring plant species, for example through the ability of foundation species to ameliorate micro-environmental conditions (Schöb *et al.* 2012; Butterfield & Callaway 2013; McIntire & Fajardo 2014). Thus, understanding the role played by microhabitats created by foundation species in sustaining biodiversity with a changing climate is crucial (Ellison *et al.* 2005; Brooker 2006), especially in Mediterranean alpine ecosystems, where facilitation by foundation species is particularly relevant for plant community diversity (Schöb, Armas & Pugnaire 2013; Michalet *et al.* 2014; Pistón *et al.* 2016). Consequently, plant–plant interactions matter, but our tests of their importance are generally pairwise. A holistic community-level approach that considers networks of interacting species would be useful and

This article is protected by copyright. All rights reserved.

might improve our understanding of species and community responses to environmental changes. However, such a network approach has hardly been employed so far (but see Verdú & Valiente-Banuet 2008; Allesina & Levine 2011; Saiz & Alados 2011).

To investigate the resistance of a community to environmental perturbations, a common approach is to perform random extinctions or target extinctions that eliminate the most and the least connected species of an interaction network first (Solé & Montoya 2001; Dunne, Williams & Martinez 2002; Memmott, Wasser & Price 2004; Verdú & Valiente-Banuet 2008). However, such an approach does not consider species sensitivity to specific environmental changes (Ives & Cardinale 2004; Curtsdotter *et al.* 2011). Functional traits can offer such a biology-informed, mechanistic link between prevailing environmental conditions and the likelihood of species survival (McGill *et al.* 2006; Garnier *et al.* 2016). Indeed, by capturing essential aspects of species' ecophysiology, functional traits determine the sensitivity of plants to biotic and abiotic factors (Grime 2001; Westoby *et al.* 2002; Ackerly 2004; Schöb *et al.* 2012; Butterfield & Callaway 2013). Consequently, traits predispose plant species to extinction under certain kinds of environmental changes (Cardinale *et al.* 2012).

In this study we present a new analytical and conceptual framework to simulate plant community persistence against environmental perturbations, showing its application to a Mediterranean alpine plant community dominated by three foundation species. We characterise the plant–plant interaction network and explore the consequences of foundation species loss for biodiversity using functional traits as a biological criterion of species' sensitivity to environmental changes. We then estimate the loss of species that is caused by the primary loss of microhabitats created by foundation species (i.e. secondary extinctions *sensu* Brodie *et al.* 2014). We hypothesised that community persistence and the rate of species loss varied among environmental change scenarios, as would the role of foundation species in modulating the species loss.

This article is protected by copyright. All rights reserved.

Materials and methods

Study area

The study was performed in the Sierra Nevada Mountains (Spain, 2725 m a.s.l., 37.08134° N, -3.38127° E), where the importance of positive plant interactions for community structure and diversity is well documented (Schöb *et al.* 2012, 2013a, b, 2014; Pistón *et al.* 2016). The site was a relatively homogeneous 0.5 ha plot, with a patchy plant community dominated by three foundation species: the cushion-forming species *Arenaria tetraquetra* spp. *amabilis* (Bory) H. Lindb. Fil. (Caryophyllaceae) and *Plantago holosteum* Scop. (Plantaginaceae), and the tussock grass *Festuca indigesta* Boiss. (Poaceae). At the closest weather station in Pradollano (2500 m a.s.l.; 37.08333 °N, -3.38333° E) mean growing season (May-September) temperature is 13.8°C (average min 6.2°C, average max 21.5°C) and mean annual precipitation is 690 mm, with a dry summer period during July and August (<http://es.climate-data.org>). The habitat belongs to the oro-Mediterranean acidic grassland type characterised by low-productivity, siliceous gravel substrate with poorly developed soils and generally low soil organic matter and water content (Schöb *et al.* 2012). The examined foundation species are known to provide positive effects on some other plant species in the study area, mainly through provision of higher soil organic matter and soil water compared to bare ground (i.e. open) areas (Schöb *et al.* 2012, 2013b; C. Schöb, unpublished data). This positive, facilitative effect was demonstrated by a better plant water status and higher reproductive output of subordinate species associated to foundation species (Schöb *et al.* 2014), indicating the dependency of subordinate species on microhabitats created by foundation species for their survival and reproduction.

Sampling was performed within the canopy area of 40 individuals of each of the three foundation species and in 10 open areas (hereafter, we refer to these samples as plots and to the each of the three foundation species and the open areas as different microhabitats, This article is protected by copyright. All rights reserved.

respectively). We identified all individuals of plant species in each plot and subsequently refer to them as subordinate plant species either occurring within the canopy of foundation species or in the open microhabitat (see Appendix S1 in Supporting Information). As plant–plant interactions are local processes acting at fine spatial scale, sampling was performed at a centimetre scale, consistent with previous studies in the study area that inferred plant interactions from spatial patterns (Schöb *et al.* 2012, 2013a, b, 2014; Pistón *et al.* 2016). Plot size was equal to the foundation species canopy size: 168 (8-406) cm² [mean (min-max)] for *Arenaria*; 149 (3-638) cm² for *Plantago*; 241 (<1-1'018) cm² for *Festuca*. In the open we sampled circular plots (27.6 cm diameter or 598 cm²) that were randomly placed in gaps between foundation species

Trait-environment relationships

We used three functional traits with a known sensitivity to environmental conditions and related to resource use strategies: specific leaf area (SLA) and leaf dry matter content (LDMC). SLA increases with soil nutrient availability (Ordoñez *et al.* 2009) and is related to photosynthesis and respiration (Garnier *et al.* 2016). LDMC increases with drought (Ackerly 2004), where physically robust species have high LDMC with high carbon-construction costs (Westoby *et al.* 2002). Along this leaf economics spectrum (Díaz *et al.* 2016), species with high LDMC and low SLA have lower photosynthetic rates and slower return on investment, indicating a conservative growth strategy and high drought stress tolerance (due to reduced water loss). Vice versa, species with high SLA and low LDMC acquire resources faster, showing high competitive ability but lower stress tolerance (i.e. higher vulnerability to stress).

Thus, we assumed that under the scenario of 1) increasing drought alone (i.e. without temperature limitation), reducing water loss through high LDMC is a relevant strategy (Grime 2001; Ackerly 2004; Liancourt *et al.* 2015); 2) combined increases of drought and temperature, LDMC may increase while SLA decreases (Liancourt *et al.* 2015); 3) increasing nutrients, a rapid resource-acquisitive strategy (i.e. high SLA) is beneficial and would confer a competitive advantage under the subsequent increase in competition (Wright *et al.* 2005; Liancourt *et al.* 2013; Garnier *et al.* 2016). A scenario with temperature increase alone was not assessed because it is reasonable to assume that in such dry alpine ecosystem temperature increase alone, without changing water availability, would result in a more drought stress (i.e. scenario 1 and 2, IPCC 2014).

The trait measurement procedure followed a standard protocol (Pérez-Harguindeguy *et al.* 2013). For each individual subordinate plant and for five individuals of each foundation species the best-developed and most healthy leaf was collected ($n = 1340$). Leaves were fully rehydrated before determining the fresh mass and the leaf area, then they were dried at 80°C for at least 72 h to determine dry mass. SLA was measured as the ratio between leaf area measured as one-sided surface area of an individual lamina (m^2) and leaf dry mass ($\text{m}^2 \text{kg}^{-1}$), and LDMC was measured as the ratio between leaf dry mass and water-saturated fresh mass (g kg^{-1}). The sampling took place at the peak of the growing season, between the end of July and the beginning of August 2011.

Plant association network

The network was built from the observed community matrix, i.e. the set of abundance data for n subordinate species growing within m microhabitats (i.e. the three foundation species and open areas), considering all plant species as nodes and co-occurrence between subordinate species and foundation species or open areas as pairwise links (Fig. 1). Then, the

This article is protected by copyright. All rights reserved.

one preferred microhabitat for each subordinate species was identified with a community structure detection algorithm, which computed optimal partitioning of species into the four microhabitats. This algorithm maximizes modularity with simulated annealing according to patterns of intra- and inter-module connections (Guimerà & Amaral 2005; Doulier & Stouffer 2015).

Species extinction model

The species extinction procedure as a consequence of simulated environmental changes consisted of three main steps. First, for each iteration of primary extinction (see below), a species trait value was sampled from its trait distribution (i.e. a normal distribution with a mean equal to each species' observed trait mean over all microhabitats and the standard deviation equal to each species' observed standard deviation). With this procedure we accounted for intraspecific trait variation at the study site (i.e. species trait plasticity, Fig. S2). Second, based on the above-mentioned relationships among leaf traits and environmental conditions, species were removed according to the following criteria. For increasing drought alone (hereafter scenario D), species were removed in order of increasing LDMC, with species having the lowest LDMC to be removed first. As increase in drought and temperature (hereafter scenario D & T) is based on both LDMC and SLA, in order to sort plant species within this bivariate trait space, a principal component analysis (PCA) was carried out, including species trait values as active variables. First, the correlation coefficients between the two variables were evaluated. Since we expected an inverse relationship between LDMC and SLA (Pérez-Harguindeguy *et al.* 2013), we selected the main axis of the PCA that described the negative correlation between these two traits. Then, species were sorted according to their coordinates on the axis and removed in order of increasing values, with species having the lowest values along the selected axis (i.e. low LDMC and high SLA) to be

This article is protected by copyright. All rights reserved.

removed first. For increasing nitrogen deposition (hereafter scenario N), species were removed in order of increasing SLA, with species having the lowest SLA to be removed first. Besides these four environmental change scenarios, we performed a random extinction model (hereafter scenario R) by randomly removing species from the network. For each scenario the corresponding trait space was divided into 25 equally spaced intervals, representing the primary extinction sequence. At each step (1, 2, 3, ..., 25), an increasing number of intervals from the trait range was removed from the viable trait space and plant species that did not fit this viable trait space were removed from the network. In other words, plant species that fell out of the range of surviving trait values were considered primary extinct. In the scenario R, an equal number of species was removed in each extinction sequence. Therefore, different to previous extinction models (Solé & Montoya 2001; Dunne *et al.* 2002; Memmott *et al.* 2004; Verdú & Valiente-Banuet 2008), such as scenario R, the biology-informed extinction sequences using functional traits could result in unequal numbers of species removed from the network during each primary extinction iteration, depending exclusively on the number of species assigned to each trait interval.

Third, besides primary extinctions, we considered also secondary extinctions (*sensu* Brodie *et al.* 2014). Since each foundation species provides specific and unique microenvironmental conditions (Ellison *et al.* 2005; Schöb *et al.* 2012, 2013; Butterfield & Callaway 2013; McIntire & Fajardo 2014), their loss can be compared to the loss of a microhabitat. Thus, we defined secondary extinctions as the loss of subordinate species due to microhabitat loss, i.e. the loss of those species that belonged to the module of a foundation species that got primary extinct.

Every primary extinction step was iterated 200 times (n replicates= 25'000). At the end of each iteration we computed: a) the proportion of surviving species (i.e. relative species richness) within the network, calculated as the sum of species that survived divided by the

This article is protected by copyright. All rights reserved.

total species richness; b) the proportion of secondary extinctions, calculated as the sum of secondary extinctions divided by the total number of extinctions (i.e. primary plus secondary); c) the persistence status of each plant species (i.e. survival, primary or secondary extinct).

Statistical analysis

We used linear mixed effects models to test the response variables (1) proportion of surviving species and (2) proportion of secondary extinctions as a function of the extinction scenarios (fixed effect), number of foundation species (fixed effect), and their interaction (fixed effect), with random effects being the iteration nested within each extinction step. To account for series autocorrelation, the steps along the sequences within each scenario were included in the first order correlation model for the residual error component (Butler *et al.* 2007). The significance of the models was tested with Wald tests.

To confirm the relevance of the model, we assessed whether subordinate species differed in their probability of surviving or getting secondary extinct by fitting a multinomial logit model with the status of the species (survived, primary or secondary extinct) as response categories and the species as predictors (Venables & Ripley 2002). The primary extinction was considered as reference level. Then, in order to explore which biotic factors make species more likely to persist or to get secondary extinct, generalised linear models (GLM) were fitted with the module membership and the abundance of each subordinate species as predictors and the likelihood (i.e. the logit estimated by the multinomial logit model) of survival or to get secondary extinct as responses. The likelihood to get primary extinct was taken as reference level for these analyses. The significance of the models was tested with ANOVA tests and likelihood-ratio χ^2 estimation.

This article is protected by copyright. All rights reserved.

Analyses were performed in R 3.1.3. (R Core Team 2015) using *igraph* for network analysis (Csárdi & Nepusz 2006), *Rnetcarto* for network modularity (Doulcier & Stouffer 2015), *asreml* for mixed effects models (Butler *et al.* 2007), *pascal* for Wald tests and *nnet* for the multinomial logit model (Venables & Ripley 2002).

Results

Plant association network response

The plant association network was composed of four modules corresponding to the four microhabitat types (Fig. 1). The *Plantago* module was the most species rich, with 11 associated subordinate species, followed by the *Arenaria* module composed of 9 associated subordinate species, the *Festuca* module with 2 associated subordinate species, and the *open* module with 3 associated subordinate species.

The three environmental change scenarios and the number of foundation species had significantly interactive effects on the proportion of surviving species (Table 1). In scenario D of increasing drought alone, in scenario D & T of increasing drought and temperature, and in scenario random, species diversity smoothly decreased in the first half of the sequence, with, on average, 50% of the species disappearing after removal of *c.* 40% of the trait space (Fig. 2). Then, in the second half of the sequence a further intensification of environmental changes caused less extinction. In scenario D, *Arenaria*, *Plantago*, and *Festuca* disappeared after removal of 39%, 23%, and 48% of the trait space, respectively, while in scenario D & T they disappeared after 23%, 26%, and 34%, respectively. These scenarios produced less secondary extinction events (4% and 5% on average, respectively) and created hump-shaped responses of secondary extinctions, which made up to 10% of the species losses when 56% and 49% of the trait spaces were removed, respectively (Fig. 3). Scenario N of increasing nitrogen deposition produced a constant steep decline in species diversity, with 50% of

species disappearing when *c.* 25% of the trait space was removed (Fig. 2). After removal of 40% of the trait space, species diversity showed a slower non-linear decrease. This scenario caused the highest secondary extinction rates (on average 20%), with a right-skewed hump-shaped pattern along the trait range removal sequence (Fig. 3). Here, *Arenaria*, *Plantago*, and *Festuca* disappeared after removal of 31%, 12% and 4% of the trait space, respectively.

Subordinate species persistence

Subordinate plant species exhibited different likelihood of survival or secondary extinction within the network. (Fig. 4, Tab. S1). Over all scenarios, module membership significantly explained the likelihood of subordinate species to survive or get secondary extinct ($G^2_3=23.17$, $P < 0.001$, $G^2_3=581.98$, $P < 0.001$, respectively; Table S2). Subordinate species associated to *Arenaria* and *Plantago* modules were significantly less likely to survive ($t=-2.58$, $P < 0.05$, $t=-2.87$, $P < 0.05$, respectively), and significantly more likely to get secondary extinct ($t=12.89$, $P < 0.001$, $t=15.21$, $P < 0.001$, respectively) compared to subordinate species associated with *Festuca* and the open microhabitat.

Discussion

Our assessment of the resistance of alpine plant communities to microhabitat loss using functional traits as a proxy for species' susceptibility showed that extinction rates were dependent on the type of environmental perturbation. Indeed, the order and timing of breakdown of the plant interaction network sustained by the foundation species differed substantially among the four environmental change drivers. The plant interaction network underlying the studied plant community was rather fragile to a simulated increase in temperature alone and increasing nitrogen deposition. Conversely, the plant interaction network resisted against moderate drought alone or drought combined with temperature

This article is protected by copyright. All rights reserved.

increases. Community persistence against these latter environmental perturbations can be explained by the fact that in these two scenarios foundation species remained longer in the network, thereby slowing down biodiversity loss compared to a random extinction simulation. These results might indicate the relevance of plant association networks in modulating the impact of different environmental changes on species diversity and community persistence. Thus, within the limits of our assumptions, predictions of the fate of species and communities depend on the knowledge of the main driver of environmental change at the location of interest and how this environmental perturbation affects the plant interaction network.

Climate warming has recently been proved to change the composition and the structure of alpine plant communities, leading to a decline and disappearance of cold-adapted plant species (Gottfried *et al.* 2012). Furthermore, Hautier *et al.* (2015) found a biodiversity decline in response to nitrogen enrichment in experimental grassland communities. Our simulation suggested that nitrogen deposition could alter plant network structure, resulting in potentially accelerated species loss. In other words, just a small shift in the corresponding functional trait space, i.e. a removal of species with lower SLA values, caused the loss of foundation species and therefore a disproportionate increase in secondary extinctions. This is consistent with a fragile ecosystem in which small environmental perturbation causes the primary extinction of a small but important fraction of species that have critical cascading effects (Fortuna & Bascompte 2006; Memmott *et al.* 2007; Rezende *et al.* 2007).

Species distribution models suggest that decreasing precipitation plays an important role in determining the potential impacts of climate change on vegetation, mainly due to habitat loss (Engler *et al.* 2011). We found that drought may not be the main driving factor for species loss in the dry alpine climate of the Sierra Nevada (Spain). We suggest that the general tolerance to drought of the species growing in this environment (Giménez-Benavides

This article is protected by copyright. All rights reserved.

et al. 2007), in particular that of foundation species, might actually allow this vegetation to better resist further increase in drought compared to other environmental perturbations.

Having shown that plant–plant interaction networks responded in different ways to different environmental perturbations, we next explored how the role of foundation species differed across the different scenarios. The emerging patterns of secondary extinctions with increasing drought alone or increasing drought combined with temperature probably resemble the pattern of extinctions following the order in species abundance, with rare species becoming extinct first (Mommott *et al.* 2004). Under these circumstances, it is expected that the loss of least connected (Solé & Montoya 2001; Dunne *et al.* 2002) or least abundant (Verdú & Valiente-Banuet 2008) species does not have important negative consequences on network structure. Conversely, in the scenarios N, where foundation species got lost early in the extinction sequence, species diversity declined rapidly also due to higher rates of secondary extinctions. This suggests that in our study system, increases in nitrogen availability might negatively affect survival of the foundation species. This might be due to increasing competitive pressure from colonizing subordinate species (Schöb *et al.* 2013a, 2014), and finally competitive exclusion of the foundation species (McAuliffe 1984).

Taken together, in this dry alpine ecosystem dominated by drought-tolerant species, moderate increases in drought stress may not have important consequences for the plant community (Miranda *et al.* 2009). This effect might particularly be due to the high drought resistance of the foundation species present (Schöb *et al.* 2012, 2013a) and the role these species play in the plant interaction network. On the other hand, changes that favour more competitive species might accelerate changes in the plant community (Michalet *et al.* 2014), also due to the extinction of the less competitive foundation species (McAuliffe 1984; Liancourt, Callaway & Michalet 2005; Maestre *et al.* 2009).

Conclusions

Our study is one of the few attempts to analyse the consequences of environmental changes on ecological networks and their species by taking into account the species' sensitivities to those environmental drivers (Tylianakis *et al.* 2008). Our simulation approach approximates species deletion according to global trait-environment relationships. Furthermore, we take into account that species are plastic and include the observed trait plasticity in our models. Thus, our trait-based extinction models in response to environmental changes are biology-informed, and therefore likely to be more ecologically meaningful than previous extinction models based only on the level of species interconnectedness (i.e. species degree) or random species removal (e.g. Solé & Montoya 2001; Dunne *et al.* 2002; Memmott *et al.* 2004; Verdú & Valiente-Banuet 2008). Nevertheless, our study has limitations in that it is mainly static: we did not account for birth and mortality changes within species, nor for range shifts of species into new potentially suitable habitats, nor the immigration of new species from lower altitudes. We did not consider potential changes in network structure nor the interaction rewiring, i.e. the establishment of new interactions. Furthermore, we ignored the potential effects of differences in microhabitat size. Nevertheless, our approach represents a conceptual advance for linking functional ecology with network theory into a unified framework that could improve predictions on community responses to environmental changes. As such, we foster its use in all types of interactions (e.g. predator–prey), environmental changes (e.g. disturbance) and corresponding response traits (e.g. body mass).

Acknowledgements

This study was financially supported by the Swiss National Science Foundation (PZ00P3_148261). We thank N. Pistón, P. Macek and F.I. Pugnaire for their help with data collection. We thank J. Bascompte, R. Michalet, M.A. Fortuna, M. Caccianiga and G.M.

This article is protected by copyright. All rights reserved.

Palamara for their fruitful discussions, and B. Schmid for helping with mixed-effects models.

In addition, we wish to thank the Associate Editor, Mikey O'Brien and Takehiro Sasaki for commenting on an earlier version of this manuscript. Authors disclose any potential sources of conflict of interest.

Data Accessibility

Network data available from the Dryad Digital Repository <http://dx.doi.org/10.5061/dryad.35q01>
R scripts uploaded as online supporting information.

References

- Ackerly, D. (2004) Functional strategies of chaparral shrubs in relation to seasonal water deficit and disturbance. *Ecological Monographs*, **74**, 25–44.
- Allesina, S. & Levine, J.M. (2011) A competitive network theory of species diversity. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5638–5642.
- Bascompte, J. & Jordano, P. (2014) *Mutualistic Networks*. Princeton University Press, Princeton and Oxford, 200 pp.
- Brodie, J.F., Aslan, C.E., Rogers, H.S., Redford, K.H., Maron, J.L., Bronstein, J.L., Groves, C.R. (2014) Secondary extinctions of biodiversity. *Trends in Ecology and Evolution*, **29**, 664–672.
- Brooker, R.W. (2006) Plant-plant interactions and environmental change: Tansley review. *New Phytologist*, **171**, 271–284.
- Butler, D., Cullis, B., Gilmour, A. & Gogel, B. (2007) Analysis of Mixed Models for S-language Environments: ASReml–R Reference Manual.
- Butterfield, B.J. & Callaway, R.M. (2013) A functional comparative approach to facilitation and its context dependence. *Functional Ecology*, **27**, 907–917.
- Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P. *et al.* (2012) Biodiversity loss and its impact on humanity. *Nature*, **489**, 326–326.
- Cavieres, L.A., Brooker, R.W., Butterfield, B.J., Cook, B.J., Kikvidze, Z., Lortie, C.J. *et al.* (2014) Facilitative plant interactions and climate simultaneously drive alpine plant

diversity. *Ecology Letters*, **17**, 193–202.

Csárdi, G. & Nepusz, T. (2006) The igraph software package for complex network research. *International Journal of Complex Systems*, 1695.

Curtsdotter, A., Binzer, A., Brose, U., de Castro, F., Ebenman, B., Eklöf, A. *et al.* (2011) Robustness to secondary extinctions: Comparing trait-based sequential deletions in static and dynamic food webs. *Basic and Applied Ecology*, **12**, 571–580.

Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S. *et al.* (2016) The global spectrum of plant form and function. *Nature*, **529**, 167–171.

Doulcier, G. & Stouffer, D. (2015) Rnetcarto: Fast Network Modularity and Roles Computation by Simulated annealing. R package version 0.2.4.

Dunne, J.A., Williams, R.J. & Martinez, N.D. (2002) Network structure and biodiversity loss in food webs: Robustness increases with connectance. *Ecology Letters*, **5**, 558–567.

Ellison, A.M., Bank, M.S., Clinton, B.D., Colburn, E.A., Elliott, K., Ford, C.R. *et al.* (2005) Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment*, **3**, 479–486.

Engler, R., Randin, C.F., Thuiller, W., Dullingers, S., Zimmermann, N.E., Araújo, M.B. *et al.* (2011) 21st century climate change threatens mountain flora unequally across Europe. *Global Change Biology*, **17**, 2330–2341.

Fortuna, M.A. & Bascompte, J. (2006) Habitat loss and the structure of plant-animal mutualistic networks. *Ecology letters*, **9**, 281–286.

Garnier, E., Navas, M. & Grigulis, K. (2016) *Plant functional diversity – Organism traits, community structure, and ecosystem properties*. Oxford University Press. 231 pp.

Giménez-Benavides, L., Escudero, A. & Iriondo, J.M. (2007) Local adaptation enhances seedling recruitment along an altitudinal gradient in a high mountain Mediterranean plant. *Annals of Botany*, **99**, 723–734.

Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Barančok, P., Alonso, J.L.B. *et al.* (2012) Continent-wide response of mountain vegetation to climate change. *Nature Climate Change*, **2**, 111–115.

Grime, J.P. (2001) *Plant strategies, vegetation processes, and ecosystem properties*, 2nd edn. Wiley, New York.

Guimerà, R. & Amaral, N.L. (2005) Functional cartography of complex metabolic networks. *Nature*, **433**, 895–900.

Hautier, Y., Tilman, D., Isbell, F., Seabloom, E.W., Borer, E.T. & Reich, P.B. (2015)

This article is protected by copyright. All rights reserved.

Anthropogenic environmental changes affect ecosystem stability via biodiversity.
Science, **348**, 336–340.

- IPCC (2014) *Climate change 2014: Impacts, Adaptations, and Bulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 688 pp.
- Ives, A.R. & Cardinale, B.J. (2004) Food-web interactions govern the resistance of communities after non-random extinctions. *Nature*, **429**, 174–177.
- Klanderud, K. (2005) Climate change effects on species interactions in an alpine plant community. *Journal of Ecology*, **93**, 127–137.
- Körner, C. (2003) *Alpine plant life. Functional plant ecology of high mountain Ecosystems*. Springer-Verlag Berlin Heidelberg. 342pp.
- Liancourt, P., Callaway, R.M. & Michalet, R. (2005) Stress tolerance abilities and competitive responses determine the outcome of biotic interactions. *Ecology*, **86**, 1611–1618.
- Liancourt, P., Spence, L.A., Song, D.S., Lkhagva, A., Sharkhuu, A., Boldgiv, B. *et al.* (2013) Plant response to climate changes varies with topography, interactions with neighbours, and ecotype. *Ecology*, **94**, 444–453.
- Liancourt, P., Boldgiv, B., Song, D.S., Spence, L.A., Helliker, B.R., Petraitis, P.S. & Casper, B.B. (2015) Leaf-trait plasticity and species vulnerability to climate change in a Mongolian steppe. *Global Change Biology*, **21**, 3489–3498.
- Losapio, G. & Schöb, C. (2017) Data from: Resistance of plant–plant networks to biodiversity loss and secondary extinctions following simulated environmental changes. Dryad Digital Repository. doi:10.5061/dryad.35q01
- Maestre, F.T., Callaway, R.M., Valladares, F. & Lortie, C.J. (2009) Refining the stress gradient hypothesis for competition and facilitation in plant communities. *Journal of Ecology*, **97**, 199–205.
- McAuliffe, J.R. (1984) Sahuaro-nurse tree associations in the Sonoran Desert: competitive effects of sahuaros. *Oecologia*, **64**, 319–321.
- McGill, B.J., Enquist, B.J., Weiher, E. & Westoby, M. (2006) Rebuilding community ecology from functional traits. *Trends in Ecology and Evolution*, **21**, 178–185.
- Memmott, J., Waser, N.M. & Price, M.V. (2004) Tolerance of pollination networks to species extinctions. *Proceedings of the Royal Society B*, **271**, 2605–2611.

- Memmott, J., Craze, P.G., Waser, N.M. & Price, M.V. (2007) Global warming and the disruption of plant-pollinator interactions. *Ecology Letters*, **10**, 710–717.
- Michalet, R., Schöb, C., Lortie, C.J., Brooker, R.W. & Callaway, R.M. (2014) Partitioning net interactions among plants along altitudinal gradients to study community responses to climate change. *Functional Ecology*, **28**, 75–86.
- Michalet, R., Maalouf, J-P., Choler, P., Clément, B., Rosebery, D., Royer J-M. *et al.* (2015) Competition, facilitation and environmental severity shape the relationship between local and regional species richness in plant communities. *Ecography*, **38**, 335–345.
- Miranda, J.D.D., Padilla, F.M., Lázaro, R. & Pugnaire, F.I. (2009) Do changes in rainfall patterns affect semiarid annual plant communities? *Journal of Vegetation Science*, **20**, 269–276.
- Ordoñez, J.C., van Bodegon, P.M., Witte, J-P.M., Wright, I.J., Reich, P.B. & Aerts, R. (2009) A global study of relationships between leaf traits, climate and soil measures of nutrient fertility. *Global Ecology and Biogeography*, **18**, 137–149.
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P. *et al.* (2013) New Handbook for standardized measurement of plant functional traits worldwide. *Australian Journal of Botany*, **61**, 167–234.
- Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N. *et al.* (2014) The biodiversity of species and their rates of extinction, distribution, and protection. *Science*, **344**, 1246752.
- Pistón, N., Schöb, C., Armas, C., Prieto I, Pugnaire, F. (2016) Contribution of co-occurring shrub species to community richness and phylogenetic diversity along an environmental gradient. *Perspectives in Plant Ecology Evolution and Systematics*, **19**, 30–39.
- R Core Team (2015) R: A Language and Environment for Statistical Computing.
- Rezende, E.L., Lavabre, J.E., Guimarães, P.R., Jordano, P. & Bascompte, J. (2007) Non-random coextinctions in phylogenetically structured mutualistic networks. *Nature*, **448**, 925–929.
- Saavedra, S., Rohr, R.P., Dakos, V. & Bascompte, J. (2013) Estimating the tolerance of species to the effects of global environmental change. *Nature Communications*, **4**, 1–6.
- Saiz, H. & Alados, C.L. (2011) Effect of *Stipa tenacissima* L. on the structure of plant co-occurrence networks in a semi-arid community. *Ecological Research*, **26**, 595–603.
- Schöb, C., Butterfield, B.J. & Pugnaire, F.I. (2012) Foundation species influence trait-based

community assembly. *New Phytologist*, **196**, 824–834.

Schöb, C., Armas, C., Guler, M., Prieto, I. & Pugnaire, F.I. (2013a) Variability in functional traits mediates plant interactions along stress gradients. *Journal of Ecology*, **101**, 753–762.

Schöb, C., Armas, C. & Pugnaire, F.I. (2013b) Direct and indirect interactions co-determine species composition in nurse plant systems. *Oikos*, **122**, 1371–1379.

Schöb, C., Prieto, I., Armas, C. & Pugnaire, F.I. (2014) Consequences of facilitation: one plant's benefit is another plant's cost. *Functional Ecology*, **28**, 500–508.

Solé, R.V. & Montoya, J.M. (2001) Complexity and fragility in ecological networks. *The Royal Society B*, **268**, 2039–2045.

Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008) Global change and species interactions in terrestrial ecosystems. *Ecology Letters*, **11**, 1351–1363.

Venables, W.N. & Ripley, B.D. (2002) *MASS: modern applied statistics with S*, Fourth edn. Springer, 495 pp.

Verdú, M. & Valiente-Banuet, A. (2008) The nested assembly of plant facilitation networks prevents species extinctions. *The American naturalist*, **172**, 751–760.

Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. (2002) Plant ecological strategies: some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*.

Wright, I.J., Reich, P.B., Cornelissen, J.H.C., Falster, D.S., Groom, P.K., Hikosaka, K. *et al.* (2005) Modulation of leaf economic traits and trait relationships by climate. *Global Ecology and Biogeography*, **14**, 411–421.

Table 1 Fixed effects test and variance component estimates (standard error) for linear mixed-effects models. The number of foundation species and its interaction with the five environmental change scenarios are predictors of species survival and secondary extinctions.

*** $P < 0.001$.

	Species survival	Secondary extinctions
Fixed effects		
Intercept	$F_{1, 13'163} = 70.0$ ***	$F_{1, 15'015} = 10.4$ ***
Scenario	$F_{3, 13'163} = 838.6$ ***	$F_{3, 15'015} = 3230.0$ ***
Foundation sp	$F_{3, 13'163} = 12310.0$ ***	$F_{3, 15'015} = 14080.0$ ***
Scenario x Foundation	$F_{9, 13'163} = 345.4$ ***	$F_{9, 15'015} = 136.2$ ***
Variance components		
Step/Iteration	9.0×10^{-3} (1.0×10^{-5})	5.3×10^{-7} (3.4×10^{-11})
Series autocorrelation		
ρ_{AR1}	0.5 (6.1×10^{-3})	0.4 (6.6×10^{-3})

Figure 1 Plant–plant network in the Sierra Nevada, Spain. Plant species are represented by nodes, whose dimensions are proportional to the species abundance. Links are proportional to the number of links (i.e. individuals) between species. Link length depends on node position according to the Kamada & Kawai force-directed algorithm (Csárdi & Nepusz, 2006).

Microhabitats are represented by rectangles, subordinate species by circles. Colours depict network modules (i.e. microhabitat preferences): blue for *Arenaria tetraquetra* spp. *amabilis*, green for *Festuca indigesta*, red for *Plantago holosteum*, and brown for open microhabitat.




Figure 2 Species diversity response (y-axis) to the four environmental change scenarios and the random model along the removal sequence (x-axis). Scenario T: increasing temperature alone; Scenario D: increasing drought alone; Scenario D & T: increasing drought and temperature; Scenario N: increasing nitrogen deposition; Scenario R: random model. Squares (*Arenaria tetraquetra* ssp. *amabilis*), circles (*Plantago holosteum*) and triangles (*Festuca indigesta*) show where each foundation species goes extinct on average. Lines fitted with a local polynomial surface (mean \pm 95% CI).




Figure 3 Secondary extinctions (y-axis) in response to the four environmental change scenarios and the random model along the removal sequence (x-axis). Scenario T: increasing temperature alone; Scenario D: increasing drought alone; Scenario D & T: increasing drought and temperature; Scenario N: increasing nitrogen deposition; Scenario R: random model. Squares (*Arenaria tetraquetra* ssp. *amabilis*), circles (*Plantago holosteum*) and triangles (*Festuca indigesta*) show where each foundation species goes extinct on average. Lines fitted with a local polynomial surface (mean \pm 95% CI).





Figure 4 Multinomial log-linear model of the differences among subordinate species (x-axis) in their probability (i.e. logit) of persistence within the network (y-axis), i.e. surviving **(a)** or getting secondary extinct **(b)**. For the 95% confidence intervals (y-axis in (b) rescaled for clarity), see Tab. S1.



Supporting information

Additional supporting information may be found in the online version of this article.

Appendix S1 Conceptual framework, trait distributions and model coefficients.

- **Figure S1** Conceptual framework.
- **Figure S2** Trait distributions.
- **Table S1** Multinomial log-linear model of the differences among subordinate species in their persistence within the network.
- **Table S2** Likelihood ratio chi-square and model coefficients for generalised linear models.

R scripts